

Connexion coronaires anormales : l'angioplastie coronaire a-t-elle une place ?

Dr Pierre AUBRY, Paris, France

Dr Philippe DEGRELL, Luxembourg, Luxembourg
pour le groupe ANOCOR



5 6 7
JUIN 2019



Liens d'intérêt potentiels : aucun



5 6 7
JUIN 2019



Connexion coronaires anormales : l'angioplastie coronaire a-t-elle une place ?



Guidelines

2018 AHA/ACC Guideline for the Management of Adults With Congenital Heart Disease: Executive Summary

| COR | LOE | Recommendations |
|-------------|------|--|
| Therapeutic | | |
| I | B-NR | <p>1. <u>Surgery</u> is recommended for AAOCA from the left sinus or AAOCA from the right sinus for symptoms or diagnostic evidence consistent with coronary ischemia attributable to the anomalous coronary artery.^{S4.4.5.2-1-S4.4.5.2-3}</p> |
| IIa | C-LD | <p>2. <u>Surgery</u> is reasonable for anomalous aortic origin of the left coronary artery from the right sinus in the absence of symptoms or ischemia.^{S4.4.5.2-4-S4.4.5.2-6}</p> |
| IIa | C-EO | <p>3. <u>Surgery</u> for AAOCA is reasonable in the setting of ventricular arrhythmias.</p> |
| IIb | B-NR | <p>4. <u>Surgery</u> or continued observation may be reasonable for asymptomatic patients with an anomalous left coronary artery arising from the right sinus or right coronary artery arising from the left sinus without ischemia or anatomic or physiological evaluation suggesting potential for compromise of coronary perfusion (eg, intramural course, fish-mouth-shaped orifice, acute angle).^{S4.4.5.2-4-S4.4.5.2-6}</p> |




Figure 5. Anomalous aortic origin of the coronary artery.

*Surgical intervention to involve unroofing or coronary revascularization for patients with concomitant fixed obstruction.

Anomalies de connexion coronaire et chirurgie

- Recommandations : ciblées pour une population jeune.
- Décisions thérapeutiques : indiquées sans tenir compte de l'âge.
- Etudes randomisées contrôlées : aucune.
- Histoire naturelle et corrigée : mal connue à long terme.
- Correction chirurgicale : techniques spécialisées.
- Echecs chirurgicaux : anévrysme, sténose cicatricielle, thrombose précoce.

Surgery for anomalous aortic origin of coronary arteries: a multicentre study from the European Congenital Heart Surgeons Association[†]

Massimo A. Padalino^{a,*}, Nicola Franchetti^a, Mark Hazekamp^b, Vladimir Sojak^b, Thierry Carrel^c,
Alessandro Frigiola^d, Mauro Lo Rito^d, Jurgen Horer^e, Regine Roussin^e, Julie Cleuziou^f, Bart Meyns^g, Jose Fragata^h,
Helena Telles^h, Anastasios C. Polimenakosⁱ, Katrien Francois^j, Altin Veshti^k, Jukka Salminen^l,
Alvaro Gonzalez Rocafort^m, Matej Nosalⁿ, Luca Vedovelli^o, Alvise Guariento^a, Vladimiro L Vida^a,
George E. Sarris^p, Giovanna Bocuzzo^q and Giovanni Stellin^a

1991-2018 = 27 years

14 centres

156 patients

0.4/patient/year/center

Table 1: Patient characteristics and preoperative presentation

| | All (%) | AORCA (%) | AOLCA (%) | Other (%) | P-value |
|-------------------------------------|----------------|----------------------------|---------------------------|---------------------------|-------------------------|
| Total patients | 156 (100) | 105 (67.3) | 35 (22.4) | 16 (10.3) | NS |
| Female | 72/156 (46.2) | 47/105 (44.8) | 18/35 (51.4) | 7/16 (43.8) | NS |
| Anomalous course of coronary artery | | | | | NS |
| Interarterial | 135/156 (86.5) | 96/105 (91.4) | 29/35 (82.9) | 10/16 (62.5) | NS |
| Intramural | 98/156 (62.8) | 77/105 (73.3) ^a | 18/35 (51.4) ^b | 3/16 (18.8) ^c | 0.022; 0.035; <0.0001 |
| Intraseptal or retroaortic | 4/156 (2.6) | 1/105 (1.0) | 2/35 (5.7) | 1/16 (6.3) | NS |
| Anterior to pulmonary artery | 1/156 (0.6) | 0/105 (0) | 0/35 (0) | 1/16 (6.3) | NS |
| Other | 16/156 (10.3) | 8/105 (7.6) | 4/35 (11.4) | 4/16 (25.0) | NS |
| Symptoms at diagnosis | 136/156 (87.2) | 94/105 (89.5) ^a | 30/35 (85.7) | 12/16 (75.0) ^b | 0.010 |
| Symptoms at effort | 103/156 (66.0) | 72/105 (68.6) | 22/35 (62.9) | 9/16 (56.3) | NS |
| Chest pain | 42/156 (26.9) | 31/105 (29.5) | 8/35 (22.9) | 3/16 (18.8) | NS |
| Cardiac arrest/low cardiac output | 21/156 (13.5) | 15/105 (14.3) | 5/35 (14.3) | 1/16 (6.6) | NS |
| Dyspnoea | 10/156 (6.4) | 4/105 (3.8) | 4/35 (11.4) | 2/16 (12.5) | NS |
| Palpitations | 7/156 (4.5) | 2/105 (1.9) | 3/35 (8.6) | 2/16 (12.5) | NS |
| Syncope | 14/156 (9.0) | 5/105 (4.8) ^a | 8/35 (22.9) ^b | 1/16 (6.3) | NS |
| Fatigue | 4/156 (2.6) | 3/105 (2.9) | 0/35 (0) | 1/16 (6.3) | NS |
| Not specified | 36/156 (23.1) | 35/105 (33.3) ^a | 1/35 (2.9) ^b | 2/16 (12.5) ^b | 0.003; <0.001 |
| No symptoms | 20/156 (12.8) | 11/105 (10.5) ^a | 5/35 (14.3) | 4/16 (25.0) ^b | 0.010 |
| Preoperative sport activity | 44/156 (28.2) | 29/105 (27.6) | 13/35 (37.1) | 2/16 (12.5) | NS |
| Age at procedure (years) | 39.5 (15-53) | 41 (19-53) ^a | 15 (12-44) ^b | 46 (16-57) | 0.070 |
| Percutaneous/CABG | 6/156 (3.8) | 6/105 (5.7) | 0/35 (0) | 0/16 (0) | NS |
| Associated cardiac disease | 33/156 (21.2) | 23/105 (21.9) | 11/35 (31.4) | 5/16 (31.3) | NS |
| Atrial septal defect | 4/156 (2.6) | 4/105 (3.8) | 0/35 (0) | 0/16 (0) | NS |
| Aortic valve anomaly | 12/156 (7.7) | 4/105 (3.8) | 5/35 (14.3) | 3/16 (18.8) | NS |
| Mitral valve anomaly | 7/156 (4.5) | 5/105 (4.8) | 2/35 (5.7) | 0/16 (0) | NS |
| Other | 10/156 (6.4) | 3/105 (2.9) | 4/35 (11.4) | 2/16 (12.5) | NS |

Numbers represent median (interquartile range) for continuous variables and n (%) for categorical variables.

Values in the same row that have different superscript letters are significantly different from each other.

AOLCA: aortic origin of left coronary artery; AORCA: aortic origin of right coronary artery; CABG: coronary artery bypass graft; NS, not significant.


Anomalies de connexion coronaire et mort subite

Annual risk of sudden cardiac death
in population 12-35 years of age
(estimation)

| | |
|----------------|---------------|
| L-ACAOS | 0.3% |
| R-ACAOS | 0.015% |


Brothers J. J Thorac Cardiovasc Surg 2010

Flow chart for ICD implantation



Age of SCD or Aborted SCD Attributed to AAOCA

NHS



Anomalous connections of coronary artery, SCD and prevalence

| CONGENITAL HEART DISEASE | SCD cases / 1 000 / year |
|---|--------------------------|
| Catecholaminergic polymorphic ventricular tachycardia | 15 |
| Hypertrophic cardiomyopathy | 10-20 |
| Brugada syndrome | 10 |
| Long QT syndrome | 5-10 |
| Idiopathic dilated cardiomyopathy | 5-10 |
| Arrhythmogenic right ventricular cardiomyopathy | 5-10 |
| Anomalous connection of left coronary artery | 3 |
| Wolf-Parkinson-White syndrome | 1 |
| Anomalous connection of right coronary artery | 0.15 |

Anomalous connections of coronary artery, SCD and prevalence

| CONGENITAL HEART DISEASE | SCD cases / 1 000 / year |
|---|--------------------------|
| Catecholaminergic polymorphic ventricular tachycardia | 15 |
| Hypertrophic cardiomyopathy | 10-20 |
| Brugada syndrome | 10 |
| Long QT syndrome | 5-10 |
| Idiopathic dilated cardiomyopathy | 5-10 |
| Arrhythmogenic right ventricular cardiomyopathy | 5-10 |
| Anomalous connection of left coronary artery | 3 |
| Wolf-Parkinson-White syndrome | 1 |
| Anomalous connection of right coronary artery | 0.15 |

| CONGENITAL HEART DISEASE in general population | cases per 1 000 |
|---|-----------------|
| Anomalous connection of right coronary artery | 3 |
| Hypertrophic cardiomyopathy | 2 |
| Wolf-Parkinson-White syndrome | 1.5 |
| Long QT syndrome | 0.5 |
| Anomalous connection of left coronary artery | 0.4 |
| Idiopathic dilated cardiomyopathy | 0.4 |
| Arrhythmogenic right ventricular cardiomyopathy | 0.4 |
| Brugada syndrome | 0.2 |
| Catecholaminergic polymorphic ventricular tachycardia | 0.1 |

Table 2: Intraoperative and postoperative data

| | All (%) | AORCA (%) | AOLCA (%) | Other (%) | P-value |
|---|---|--------------------------------|-------------------------------|--------------------------------|---------------------|
| Type of surgical procedure* | Median age at procedure* | | | | |
| Unroofing ^a | 88/156 (56.4)32 years (IQR 14–50; range 0.5–67) | 66/105 (62.9) | 19/35 (54.3) | 3/16 (18.8) | NS |
| Coronary reimplantation ^a | 30/156 (19.2)37 years (IQR 16.5–47; range 0.5–64) | 26/105 (24.8) | 2/35 (5.7) | 2/16 (12.5) | NS |
| CABG ^b | 24/156 (15.4)58.5 years (IQR 47–66; range 1–70) | 8/105 (7.6) | 7/35 (20.0) | 9/16 (56.3) | NS |
| Other ^a | 14/156 (9.0)32.5 years (IQR 15–43; range 3–63) | 5/105 (4.8) | 7/35 (20.0) | 2/16 (12.5) | NS |
| CPB time (min) | 73 (54–103) | 68 (52–91) | 84 (68–116) | 88 (47–137) | NS |
| CC time (min) | 47 (35–69) | 43 (34–66) | 53 (41–70) | 58 (37–78) | NS |
| ICU (days) | 1 (1–2) | 1 (1–2) | 1 (1–2) | 2 (1–3) | NS |
| Major postoperative complications | | | | | |
| Low cardiac output syndrome | 9/156 (5.8) | 2/105 (1.9) ^a | 5/35 (14.3) ^b | 2/16 (12.5) | 0.010 |
| Early reintervention | 7/156 (4.5) | 3/105 (2.9) | 3/35 (8.6) | 1/16 (6.3) | NS |
| Right coronary stenting | 2/156 (1.3) | 2/105 (1.9) | 0/35 (0) | 0/16 (0) | NS |
| Balloon dilatation of pulmonary stenosis | 1/156 (0.6) | 0/105 (0) | 1/35 (2.9) | 0/16 (0) | NS |
| Removal of thrombus from left main coronary | 1/156 (0.6) | 0/105 (0) | 1/35 (2.9) | 0/16 (0) | NS |
| Coronary artery bypass graft | 1/156 (0.6) | 1/105 (1.0) | 0/35 (0) | 0/16 (0) | NS |
| Heart transplant | 1/156 (0.6) | 0/105 (0) | 1/35 (2.9) | 0/16 (0) | NS |
| Sinus Valsalva aneurysm resection | 1/156 (0.6) | 1/105 (1.0) | 0/35 (0) | 0/16 (0) | NS |
| Other | 1/156 (0.6) | 0/105 (0) | 0/35 (0) | 1/16 (6.3) | NS |
| Mechanical support | 6/156 (3.8) | 1/105 (1.0)^a | 3/35 (8.6)^b | 2/16 (12.5)^b | 0.050; 0.050 |
| ECMO | 2/156 (1.3) | 0/105 (0) | 1/35 (2.9) | 1/16 (6.3) | NS |
| Intra-aortic balloon pump | 2/156 (1.3) | 0/105 (0) | 1/35 (2.9) | 1/16 (6.3) | NS |
| Ventricular assist device | 1/156 (0.6) | 0/105 (0) | 1/35 (2.9) | 0/16 (0) | NS |
| Impella device | 1/156 (0.6) | 1/105 (1.0) | 0/35 (0) | 0/16 (0) | NS |
| Hospital deaths | 2/156 (1.3) | 0/105 (0) | 1/35 (2.9) | 1/16 (6.3) | NS |
| Minor postoperative complications | | | | | |
| Pericardial/pleural effusion | 6/156 (3.8) | 5/105 (4.8) | 0/35 (0) | 1/16 (6.3) | NS |
| Arrhythmia | 4/156 (2.6) | 2/105 (1.9) | 1/35 (2.9) | 1/16 (6.3) | NS |
| Sepsis/infection | 3/156 (1.9) | 3/105 (2.9) | 0/35 (0) | 0/16 (0) | NS |
| Respiratory insufficiency | 2/156 (1.3) | 1/105 (1.0) | 1/35 (2.9) | 0/16 (0) | NS |
| Aortic insufficiency (mild-moderate) | 2/156 (1.3) | 2/105 (1.9) | 0/35 (0) | 0/16 (0) | NS |

Activité chirurgicale 2017 en France



ATIH
AGENCE TECHNIQUE
DE L'INFORMATION
SUR L'HOSPITALISATION



| Base Nationale Publique et Privée - 2018 (mise à jour hebdomadaire) Répartition des GHM pour l'acte CCAM | | | | |
|--|--------|--|----------|-------|
| DDEA001 : Réimplantation d'une artère coronaire pour anomalie congénitale d'origine, par thoracotomie avec CEC | | | | |
| CMD | GHM | Libellé | Effectif | DMS |
| 05 | 05C062 | Autres interventions cardiothoraciques, âge supérieur à 1 an, ou vasculaires quel que soit l'âge, avec circulation extracorporelle, niveau 2 | 29 | 8,97 |
| 05 | 05C061 | Autres interventions cardiothoraciques, âge supérieur à 1 an, ou vasculaires quel que soit l'âge, avec circulation extracorporelle, niveau 1 | 19 | 8,89 |
| 05 | 05C074 | Autres interventions cardiothoraciques, âge inférieur à 2 ans, avec circulation extracorporelle, niveau 4 | 9 | 36,11 |
| 05 | 05C064 | Autres interventions cardiothoraciques, âge supérieur à 1 an, ou vasculaires quel que soit l'âge, avec circulation extracorporelle, niveau 4 | 6 | 33,83 |
| 05 | 05C073 | Autres interventions cardiothoraciques, âge inférieur à 2 ans, avec circulation extracorporelle, niveau 3 | 4 | 18,25 |
| 05 | 05C063 | Autres interventions cardiothoraciques, âge supérieur à 1 an, ou vasculaires quel que soit l'âge, avec circulation extracorporelle, niveau 3 | 4 | 18,00 |
| 05 | 05C072 | Autres interventions cardiothoraciques, âge inférieur à 2 ans, avec circulation extracorporelle, niveau 2 | 1 | 15,00 |
| 05 | 05C071 | Autres interventions cardiothoraciques, âge inférieur à 2 ans, avec circulation extracorporelle, niveau 1 | 1 | 7,00 |
| | | | | 73 |

5 6 7
JUIN 2019



Connexion coronaires anormales : l'angioplastie coronaire a-t-elle une place ?

Anomalous connections of coronary artery and PCI

Six-Month Success of Intracoronary Stenting for Anomalous Coronary Arteries Associated With Myocardial Ischemia

- First and short series of 14 patients
- Objective evidence of ischemia
- 9 ARCA with interarterial course
- 44-72 year-old.
- Bare-metal stents (BMS)
- No procedural complications
- Resolution of myocardial ischemia on stress testing at follow-up

Doorey et al. Am J Cardiol 2000

Anomalous connections of coronary artery and PCI

Six-Month Success of Intracoronary Stenting for Anomalous Coronary Arteries Associated With Myocardial Ischemia

- First and short series of 14 patients
- 44-72 year-old
- 9 ARCA with interarterial course
- Objective evidence of ischemia
- Bare-metal stents (BMS)
- No procedural complications
- Resolution of myocardial ischemia on stress testing at follow-up

Doorey AJ et al. Am J Cardiol 2000

Origin of the Right Coronary Artery from the Opposite Sinus of Valsalva in Adults: Characterization by Intravascular Ultrasonography at Baseline and After Stent Angioplasty

- 42 patients with ARCA and interarterial intramural proximal course
- Mean age 48 ± 12 years (12-73)
- Symptomatic, positive stress test, VUS surface reduction >50%)
- IVUS-guided PCI
- Successful PCI in all patients with 93% of drug-eluting stents (DES)
- Improved symptoms at one-year follow-up (30 patients)
- 13% restenosis rate at 5-year follow-up
- No ACAOS-related deaths during follow-up

Angelini P et al. Cathet Cardio Interv 2015



5 6 7
JUIN 2019



Connexion coronaires anormales : l'angioplastie coronaire a-t-elle une place ?




Connexions coronaires anormales et angioplastie





5 6 7
JUIN 2019

APPA
Ensemble, imaginons la cardiologie de demain


Connexions coronaires anormales et angioplastie



Connexions coronaires anormales et angioplastie



Scanner - 6 mois



5 6 7
JUIN 2019



Connexion coronaires anormales : l'angioplastie coronaire a-t-elle une place ?



ANOCOR stenting registry

- ARCA with interarterial course with/without intramural pathway
- Age \geq 25 years
- No history of aborted sudden death
- Ischemic symptoms or documented myocardial ischemia
- No significant associated right CAD
- IVUS/OCT guidance
- 6-12 month CT follow-up
- Clinical follow-up at 6, 12 and 60 months

ANOCOR stenting registry

- ARCA with interarterial course with/without intramural pathway
- Age \geq 25 years
- No history of aborted sudden death
- Ischemic symptoms or documented myocardial ischemia
- No significant associated right CAD
- IVUS/OCT guidance
- 6-12 month CT follow-up
- Clinical follow-up at 6, 12 and 60 months

ANOCOR stenting registry

Baseline and angiographic characteristics

| Inclusion period: 2014-2018 | n=16 |
|------------------------------------|------------|
| Mean age (years) | 52 (29-81) |
| Male sex (%) | 10 (63) |
| Presentation | |
| ACS (%) | 2 (12.5) |
| Stable angina (%) | 11 (69) |
| Silent ischemia (%) | 2 (12.5) |
| Syncope (%) | 1 (6) |
| Angiography | |
| ARCA with interarterial course (%) | 16 (100) |
| Intramural segment (%) | 11 (69) |

ANOCOR stenting registry

Procedural characteristics

| Procedural characteristics | n=16 |
|------------------------------|----------|
| Successful stenting (%) | 16 (100) |
| DES use (%) | 15 (94) |
| Radial access (%) | 7 (44) |
| IVUS/OCT guidance (%) | 13 (82) |
| Mean stent diameter (mm) | 3.4 |
| Mean stent length (mm) | 25 |
| Mean fluoroscopic time (min) | 19 |

ANOCOR stenting registry

Outcomes

| Outcomes | n=16 |
|-----------------------------------|-----------|
| Mean troponin (microg/L) at day 1 | 0.51 |
| In-hospital complications (%) | 0 (0) |
| Mean follow-up (months) | 27 (1-60) |
| In-stent restenosis rate (%) (M8) | 2 (12.5) |
| Stent compression on CT-scan (%) | 0 (0) |
| Death during follow-up (%) | 0 (0) |

Conclusions

- Correction chirurgicale d'une connexion coronaire anormale : doit faire ses preuves dans certains cas.
- Stenting d'une connexion coronaire droite anormale : possible sans risques majeurs.
- Technique interventionnelle intégrable dans l'algorithme décisionnel : besoin d'une cohorte et d'un suivi conséquents.
- Connexion coronaire droite anormale avec symptomatologie d'allure ischémique chez un patient >30 ans : pensez à l'angioplastie.

5 6 7
JUIN 2019



Staff ANOCOR

Contact : PCAUBRY@YAHOO.FR

